

Quark droplets stability induced by external magnetic field

D. Ebert*

Institut für Physik, Humboldt-Universität zu Berlin, 10115 Berlin, Germany

K. G. Klimenko†

Institute of High Energy Physics, 142281, Protvino, Moscow Region, Russia

The influence of a constant homogeneous external magnetic field H on the formation and stability of quark droplets is investigated within a simple Nambu – Jona-Lasinio model by using a thermodynamic approach. For a vanishing magnetic field stable quark droplets, which are schematically the bags of massless quarks, are allowed to exist only at $G > G_{bag}$, where G is the quark coupling constant, $G_{bag} = 1.37G_{crit}$, and G_{crit} is the value of the coupling constant above which chiral symmetry is spontaneously broken down. On the other hand, a nonvanishing external magnetic field can induce the stability of quark droplets so that they may exist even at $G < G_{bag}$. In this case, depending on the value of H , quark droplets are composed either of massive or massless quarks.

PACS numbers: 11.30.Qc, 12.39.Fe, 26.60.+c

Keywords: Nambu – Jona-Lasinio model; chiral phase transitions; quark matter stability; magnetic catalysis effect

I. INTRODUCTION

At present time, it is generally accepted that QCD is the theory of strong interactions between mesons and baryons. Moreover, it should describe the physical phenomena in dense hadronic matter as well. Since the usual (perturbative) methods of QCD are applicable only at sufficiently high values of the baryon density ρ , in the region of low and moderate densities one should use nonperturbative methods, e.g. lattice QCD calculations [1], or effective field theory considerations. The latter ones are especially useful if the influence of external magnetic field on the properties of dense hadronic matter is investigated (in this case the incorporation of strong magnetic field into lattice calculations is not elaborated sufficiently). It is well-known that at low (baryon) densities $\rho < \rho_0$, where $\rho_0 = 0.16 \text{ fm}^{-3}$ is the normal density of baryons inside nuclei, hadronic matter is just the ordinary nuclear matter. The widely-known relativistic effective theory for nuclei and nuclear matter is the Walecka model [2, 3]. In this model the protons and neutrons, i.e. the simplest baryons (nucleons), are point particles. Moreover, the existence of stable objects with large baryonic number, which are droplets of simplest massive baryons and hence might be identified with nuclei, is predicted in the framework of it with the help of the condensed matter (thermodynamic) approach.

Unfortunately, the Walecka model is unrealistic in the region of moderate densities near to the chiral phase transition, where it is necessary to take into account the internal quark structure of protons and neutrons. Similar conditions exist in the experiments on relativistic ion-ion collisions, where it is expected that for a short time stable clumps of quark matter (usually called quark droplets) with rather large baryonic number might be created. Another example is presented by neutron stars, where under extreme pressure nucleons might fuse together, forming droplets of quarks (evidently, the baryonic density inside quark droplets is larger, than ρ_0).

One of the widely-known effective theory of quark-hadronic matter is the Nambu – Jona-Lasinio (NJL) model [4], which was successfully used in describing low-energy meson physics [5], diquarks [6, 7] and meson-baryon interactions [8, 9]. In the framework of the NJL model quarks are treated as elementary point-like particles, whereas mesons are collective quark-antiquark excitations of its ground state. This model can also serve as a starting point for investigations of the mass spectrum of the octet and decuplet of baryons (see, e.g. [10, 11, 12]). However, in order to be applicable to dense quark-hadronic matter, the NJL model should predict the existence of stable multi-quark droplets as well, i.e. of stable finite-size objects in space, which are composed of more than 3 quarks and do not tend to shrink to a point or to expand over the whole space. One particular way to investigate such objects in the NJL model and which we will follow in this paper is based on the use of the general condensed matter approach restricted to vanishing temperature T . In this case a stable multi-quark object of the system is identified with a droplet of the dense quark phase, surrounded by the vacuum phase with zero baryonic density. Hence, if there is the thermally equilibrated coexistence of these two phases, then one may conclude about the existence of stable quark droplets. Just in this way, the problem of quark matter stability was investigated at zero external magnetic field H in [13, 14, 15] in the framework of a chirally symmetric NJL model. Let us quote the two main results of these papers: i) at $H = 0$, stabilization of (multi)quark

*Electronic address: debert@physik.hu-berlin.de

†Electronic address: kklm@mx.ihep.su

droplets is only possible, when they consist of massless quarks, ii) the stability requirement puts strong constraints on the acceptable values of model parameters (coupling constant G , cutoff parameter Λ etc). In particular, it was shown that in the thermodynamic approach based on the NJL model, quark droplets can be schematically interpreted as MIT-type bags. However, in the MIT 3-quark bag model the bag constant is a phenomenological input parameter [16], whereas in the NJL model it is a dynamical quantity originating from the dynamical breaking of chiral symmetry.

Taking into account the fact that inside neutron stars the magnetic field strength might reach extremely high values (the influence of ultra-strong magnetic fields on physical processes inside neutron stars and on properties of dense quark matter are under permanent investigations [17, 18, 19, 20]), it seems to us important to reinvestigate the problem of the quark droplet stability in the framework of NJL-type models at $H \neq 0$. In the present paper, we study, in particular, the influence of a constant homogeneous external magnetic field H in a simple NJL model with two quark flavors. In Section 2, the stability constraints on the model parameters as well as on the chemical potential μ are presented. Apart from this, the phase structure of the model is found at $T = 0$, $H = 0$ and $\mu \neq 0$. On this basis, it is shown that at $H = 0$ stability of multi-quark droplets, which consist here only of massless quarks, occurs at $G > G_{bag} \equiv 1.37G_{crit}$ with $G_{crit} = \pi^2/(3N_f\Lambda^2)$ and N_f being the number of flavored quarks. In Sections 3, 4 and 5 the problem of the existence of multi-quark droplets is discussed at $H \neq 0$. In particular, it is argued on the basis of our earlier papers [21, 22] that for each value G smaller than G_{bag} , the stability of quark droplets is induced by an external magnetic field. Moreover, for each $G \in (G_{crit}, G_{bag})$ it is shown that in contrast to the case $H = 0$ [13], there exists an external magnetic field value $H_\alpha \sim 10^{19}$ Gauss such, that at $0 < H < H_\alpha$ multi-quark droplets may be formed from massive quarks. Moreover, we have shown using energetical arguments that the magnetic field promotes the creation of stable quark droplets and enhances the stability of it even at $G > G_{bag}$.

II. QUARK DROPLET STABILIZATION AT $H = 0$

A. NJL model and stability constraint

Let us first give several (very approximative) arguments motivating the chosen structure of our QCD-motivated NJL model introduced below. For this aim, consider two-flavor QCD, $N_f = 2$, with nonzero chemical potential and color group $SU_c(N_c)$ (in our case $N_c = 3$). By integrating in the generating functional of QCD over gluons and further “approximating” the nonperturbative gluon propagator by a δ -function, one arrives at an effective local chiral four-quark interaction of the NJL type describing low energy hadron physics. Finally, by performing a Fierz transformation of the interaction term and taking into account only scalar and pseudo-scalar ($\bar{q}q$) interactions, one obtains a four-fermionic model given by the following Lagrangian (in Minkowski space-time notation) ¹

$$L = \bar{q}[i\gamma^\nu\partial_\nu + \mu\gamma^0]q + G[(\bar{q}q)^2 + (\bar{q}i\gamma^5\bar{\tau}q)^2]. \quad (1)$$

In (1) $\mu > 0$ is the chemical potential of quarks which is the same for all flavors. Furthermore, we use the notations $\bar{\tau} \equiv (\tau^1, \tau^2, \tau^3)$ for Pauli matrices in the flavor space. Clearly, the model (1) is invariant under the continuous chiral $SU(2)_L \times SU(2)_R$ group. Apart from this, it is invariant under the discrete chiral transformation, $q \rightarrow i\gamma^5 q$. (Since the baryonic charge is conserved, there exists also a corresponding $U(1)_V$ -invariance of the model). At tree level, the Lagrangian (1) contains besides of the chemical potential only one free model parameter, i.e. the coupling constant G . Clearly, when including quantum effects (loops) and performing necessary regularizations, an additional cutoff parameter Λ will appear.

An illustrative way for the investigation of stable quark droplets in quantum field theory is based on a schematic analogy with the thermodynamical approach to the liquid-gas first order phase transition. In such a picture one can consider the quark droplet as a droplet of the dense liquid phase of the system, whereas the low-density gas phase corresponds to the vacuum (the zero density state) which surrounds the quark droplet (strictly speaking, such an analogy makes really sense only for systems at finite temperature). Moreover, we suppose that the two phases are in the thermal and chemical equilibrium. Of course, in this way, i.e. when ignoring surface effects of droplets, we have an adequate consideration of the quark matter stability problem only for quark droplets which contain a very large number of quarks inside a large volume (for comparison see [23], where surface effects of droplets were taken into account). It turns out that not for all values of input parameters of the system this schematic picture of the existence of stable quark droplets is realized. Thus, the stability problem of quark droplets arises, which might be expressed in terms of several quantitative constraints on the parameters of the system. To solve these constraints means to find the values of the parameters of the system at which there might appear stable quark droplets at moderate densities. Let us next describe these stability constraints.

¹ The most general Fierz transformed four-fermion interaction would include additional vector and axial-vector ($\bar{q}q$) as well as scalar and pseudo-scalar, vector and axial-vector-like (qq)-interactions. However, in order to simplify the studying of the quark matter stability problem in the presence of external magnetic field, these additional terms are omitted. The role of the interaction in the diquark channel is discussed in the last Section 5.

Recall first that in the thermal and chemical equilibrium between vacuum and dense quark matter phases their (critical) chemical potentials, denoted as μ_c , are equal. Then, from the analogy with condensed matter physics, one can get the following relations between thermodynamical and dynamical parameters at which the two phases do coexist (called below stability constraints):

$$m_{dense} < \mu_c < m_{vac}. \quad (2)$$

Here m_{vac} is the dynamical mass of one-quark excitations of the vacuum and m_{dense} is the mass of quarks in the dense phase, i.e. inside the quark droplets (see also [13]).² Note that the values of $m_{dense}, \mu_c, m_{vac}$ are supplied by the thermodynamic potential $\Omega(m; \mu)$ of the system, where the average of the composite scalar σ -field $m = \langle \sigma \rangle$ is the order parameter of spontaneous chiral symmetry breaking which in the thermal equilibrium is the global minimum point m_o of the function $\Omega(m; \mu)$ versus m . Clearly, m_o equals to the dynamical quark mass which is determined by the gap equation $m_o = -2G < \bar{q}q >$. (Obviously, if $m_o = 0$, the ground state is chirally invariant.) Hence, in order to find all the admissible values of parameters of the model, at which stable quark droplets could exist in the quantum system at moderate density, it is sufficient to solve the stability constraint (2) with respect to these parameters. On this route one should perform several technical steps: i) get the thermodynamic potential $\Omega(m; \mu)$; ii) find the global minimum point of $\Omega(m; \mu)$ vs m and establish its dependence on the external parameters of the theory thereby obtaining the phase structure of the model; iii) choose all the first order phase transition critical points μ_c obeying the constraint (2).³ The corresponding values of allowed parameters obtained in this way are just the physically acceptable ones, at which stable quark matter might be described by the initial model.

Now let us carry out this scheme on the basis of the chosen NJL model.

B. Solution of the stability constraint at $H = 0$

In the mean field approximation, the thermodynamic potential of the model (1) is given by the expression [13, 22, 24]:

$$\Omega(m; \mu) = \frac{m^2}{4G} - 2N_c N_f \int \frac{d^3 p}{(2\pi)^3} \left\{ E_p + \theta(\mu - E_p)(\mu - E_p) \right\}, \quad (3)$$

where $E_p = \sqrt{m^2 + \vec{p}^2}$, N_c is the number of colors, i.e. $N_c = 3$, and $N_f = 2$. Using for the divergent integral in (3) a Lorentz noninvariant regularization (this procedure is simply the replacement of the infinite integration region in (3) in favour of the compact one: $\vec{p}^2 \leq \Lambda^2$, where the cutoff parameter Λ is an additional parameter of the theory to be fitted by the experimental data), one can find the following expression

$$\begin{aligned} \Omega(m; \mu) = & \frac{m^2}{4G} - \frac{3N_f}{8\pi^2} \left[\Lambda(2\Lambda^2 + m^2)\sqrt{m^2 + \Lambda^2} - m^4 \ln \left(\frac{\Lambda + \sqrt{m^2 + \Lambda^2}}{m} \right) \right] - \\ & - \frac{N_f \theta(\mu - m)}{8\pi^2} \left[\mu(2\mu^2 - 5m^2)\sqrt{\mu^2 - m^2} + 3m^4 \ln \left(\frac{\mu + \sqrt{\mu^2 - m^2}}{m} \right) \right]. \end{aligned} \quad (4)$$

1. The vacuum case $\mu = 0$.

In this case, from (4) one can get the following stationarity equation:

$$\frac{\partial}{\partial m} \Omega(m; 0) = \frac{3N_f m}{2\pi^2} \left\{ \frac{\pi^2}{3N_f G} - F(m, \Lambda) \right\} = 0, \quad (5)$$

where

$$F(m, \Lambda) = \Lambda \sqrt{m^2 + \Lambda^2} - m^2 \ln \left(\frac{\Lambda + \sqrt{m^2 + \Lambda^2}}{m} \right). \quad (6)$$

² Indeed, one can interpret the chemical potential of any fermionic system as the lowest energy, needed for the particle in order to abandon the system. The equilibrium value μ_c must be smaller than the mass of the vacuum one-particle excitations m_{vac} (such particles have enough energy for leaving the vacuum, which is empty due to this reason). However, in order that quarks do not leave the dense phase, their mass m_{dense} must be smaller than μ_c – the minimum particle energy, necessary for leaving the system.

³ At the critical line μ_c of the 1st order phase transition, the order parameter m_o changes its value by a jump, i.e. at this line it is a discontinuous function vs input parameters of the model. At the critical line of the second order phase transition the order parameter is a continuous function, but first- or higher order derivatives of it are discontinuous.

The function $F(m, \Lambda)$ monotonically decreases on the interval $m \in (0, \infty)$ such that $F(0, \Lambda) = \Lambda^2$ and $F(\infty, \Lambda) = 0$ ($F(m, \Lambda) = 2\Lambda^3/(3m) + o(\Lambda/m)$ at $m \rightarrow \infty$). Hence, equation (5) has a nontrivial solution $m_o(G, \Lambda) \equiv M \neq 0$ if and only if the constraint $\frac{\pi^2}{3N_f G} - F(0, \Lambda) < 0$ (or $G \geq G_{crit} \equiv \frac{\pi^2}{3N_f \Lambda^2}$) is fulfilled. In this case, the global minimum of the function $\Omega(m; 0)$ lies at M . Thus, both the continuous and discrete chiral symmetries of the NJL-model are spontaneously broken. (At $G \leq G_{crit}$ the stationarity equation (5) has only the trivial solution $m = 0$. Thus, in this case the vacuum is chirally invariant, and fermions have zero mass. Since this phase is not adequate to the real QCD vacuum at $T = \mu = 0$, we will not consider in detail the undercritical values of the coupling constant in the present paper.) Evidently, among the three parameters Λ, G, M only two are independent ones at $G \geq G_{crit}$. In the following it is sometimes convenient to use Λ and M as independent free parameters of the model at $G \geq G_{crit}$. Clearly, M is here the mass of the one-quark excitation of the chirally-noninvariant vacuum with zero baryon density which, at $H = 0$, is just the quantity m_{vac} from (2). Then, $G = \frac{\pi^2}{3N_f F(M, \Lambda)}$ and there is a one-to-one correspondence between $G > G_{crit}$ and $M > 0$.

2. The case $\mu \neq 0$.

Note first of all that for $G \leq G_{crit}$ and arbitrary values of μ , the chiral symmetry is not spontaneously broken down. However, above the critical coupling constant the situation is more complicated. Recall, that in this case the phase structure of the simplest NJL model with $N_f = 1$ and Lorentz invariant regularization was investigated in [25]. Using the same methods, it is easy to obtain the phase structure of model (1) for the Lorentz-noninvariant regularization considered above. Thus, without any preliminary calculational discussions, we present the phase portrait of the NJL model (1) schematically on Fig. 1. On this figure one can see the (M, μ) -plane, which is divided into three domains: A, B, C. There is a symmetric dense phase with massless quarks for points $(M, \mu) \in A$. In the region B, the phase with zero baryon density is realized. In this phase the chiral symmetry is spontaneously broken down, and quarks acquire a mass M . (In this sense, phase B models the real QCD vacuum.) Finally, the region C on this figure corresponds to the dense chirally noninvariant phase, in which quarks have a nonzero μ -dependent mass. The boundaries between these phases are the critical lines of second order phase transitions (solid lines on Fig. 1), or first order ones (dashed curves on Fig. 1). In particular, the curve $l_{BC} = \{(M, \mu) : \mu = M\}$ is the critical line of the second order phase transitions, since the second order derivative $\partial^2 m_o / (\partial \mu)^2$ of the global minimum point m_o of the potential $\Omega(m; \mu)$ is a discontinuous function vs μ on this curve. The curve $l_{AB} = \{(M, \mu) : \mu = \mu_c(M)\}$ (all points of this critical curve lie below the line $\mu = M$) is defined by the equation

$$\Omega(M; \mu) = \Omega(0; \mu), \quad (7)$$

i.e. on this line the order parameter of the system – the global minimum point m_o – changes its value by a jump from M to 0, and a first order phase transition takes place. Solving (7) in the region $\mu \leq M$, one can obtain the manifest form of this curve:

$$\mu_c(M) = \left\{ 3\Lambda^3 \sqrt{M^2 + \Lambda^2} - 3\Lambda^4 - \frac{3}{2}M^2 F(M, \Lambda) \right\}^{1/4}, \quad \text{where } M \geq M_c = 0.56\Lambda. \quad (8)$$

Moreover, there are two tricritical points α and β on this phase portrait.⁴ Their coordinates are: $\alpha = (\tilde{M}, \tilde{\mu})$, $\beta = (M_c, M_c)$, where $\tilde{M} = 0.31\Lambda$, $\tilde{\mu} = 0.37\Lambda$ and M_c is defined in (8).

Now one can easily see that constraints (2) are fulfilled at $M \geq M_c$ (or in terms of coupling constants G and G_{crit} , stability constraints (2) are valid only at $G > G_{bag} \equiv 1.37G_{crit}$). Indeed, for each such value of M there is a corresponding critical value of the chemical potential $\mu_c(M)$ (8) (recall, that $0 < \mu_c(M) < M$) at which the two phases do coexist. In other words, in this case stable droplets of the dense phase A are surrounded by the chirally noninvariant vacuum (phase B) with zero baryon density. Obviously, the inequalities (2) are valid, since in the case under consideration $m_{dense} = 0$ and $m_{vac} = M$.⁵ The baryon density inside the quark droplet is easily calculated from the relation:

$$\rho_{drop}(M) = -\frac{\partial \Omega(0; \mu)}{N_c \partial \mu} \Big|_{\mu=\mu_c(M)} = \frac{N_f}{3\pi^2} \mu_c^3(M). \quad (9)$$

⁴ A point of the phase diagram is called a tricritical one if, in an arbitrary small vicinity of it, there are first- as well as second-order phase transitions.

⁵ It might appear that points of the 1st order phase transitions critical line, which connects two tricritical points α and β of Fig. 1, are also suitable ones. However, in this case there is the coexistence of two dense phases, and the physical vacuum cannot be identified with one of them. So, we reject the corresponding values of $M \in (M, M_c)$ at $H = 0$.

Now, in the framework of the NJL model, one can imagine for $M \geq M_c$ the following quark matter transition in dependence on the average baryonic density ρ : At sufficiently small ρ a rather small amount of quark droplets, having a rather small size, appears in the space. The droplets are composed of massless quarks and surrounded by the vacuum with zero baryon density; however, inside each droplet the baryonic density equals to $\rho_{drop}(M)$ (9). With growth of ρ the number of droplets as well as their sizes are increasing. When ρ reaches the value $\rho_{drop}(M)$, the total space is occupied by massless quarks.

Finally, notice that the existence of stable quark droplets surrounded by a chirally noninvariant vacuum was also considered in the framework of a NJL model in [13]. However, there the problem was investigated only for three physically interesting sets of parameters: $M = M_1 \equiv 0.48\Lambda$, $\Lambda = 650$ MeV (set 1), $M = M_2 \equiv 0.67\Lambda$, $\Lambda = 600$ MeV (set 2) and $M = M_3 \equiv 0.88\Lambda$, $\Lambda = 570$ MeV (set 3). Of course, the corresponding results are reproduced by our ones: for set 1 stable quark droplets are forbidden, but for sets 2,3 stable droplets are formed from massless quarks. In particular, it follows from (9) that $\rho_{drop} = 2.86\rho_o$ for set 2 and $\rho_{drop} = 4.19\rho_o$ for set 3 (here ρ_o is the ordinary nuclear matter density). In the present paper we have generalized these considerations and found the exact interval $M \geq 0.56\Lambda \equiv M_c$ corresponding to stable quark droplet formation.

III. PHASE STRUCTURE AT $H \neq 0$

Let us next study the phase structure of the NJL model (1) containing u - and d -quarks with electric charges $e_1 = 2|e|/3$ and $e_2 = -|e|/3$ (e is the electric charge of electrons), respectively. Due to the difference in quark electric charges, the continuous chiral symmetry is violated in the presence of an external magnetic field, but the discrete one, $q \rightarrow i\gamma^5 q$, is not. It is useful to consider first the vacuum case with $\mu = 0$, where the thermodynamic potential of the system looks like:

$$\Omega(m; 0, H) = \frac{m^2}{4G} + \sum_{i=1}^2 \frac{N_c |e_i H|}{8\pi^2} \int_0^\infty \frac{ds}{s^2} \exp(-sm^2) \coth(|e_i H|s), \quad (10)$$

where $N_c = 3$. This expression can easily be transformed into the following one (see, e.g., [22]):

$$\Omega(m; 0, H) = \Omega(m; 0) - \sum_{i=1}^2 \frac{3(e_i H)^2}{2\pi^2} \left\{ \zeta'(-1, x_i) - \frac{1}{2}[x_i^2 - x_i] \ln x_i + \frac{x_i^2}{4} \right\}, \quad (11)$$

where $\Omega(m; 0)$ is given in (4) at $\mu = 0$ and $x_i = m^2/(2|e_i H|)$. Moreover, $\zeta'(-1, x) = d\zeta(\nu, x)/d\nu|_{\nu=-1}$, where $\zeta(\nu, x)$ is the generalized Riemann zeta-function [26]. The thermodynamic potential (10)-(11) is an even function with respect to m . This fact reflects the symmetry of the original model under discrete chiral transformation $q \rightarrow i\gamma^5 q$ in the presence of an external magnetic field. Obviously, this symmetry might be spontaneously broken if the global minimum point m_o of $\Omega(m; 0, H)$ is unequal to zero, but remains intact if $m_o = 0$. In the case $G > G_{crit}$, the global minimum point of $\Omega(m; 0, H)$ is the solution of the stationarity equation (see [22]):

$$\frac{2\pi^2}{3} \frac{\partial}{\partial m} \Omega(m; 0, H) = m \{ 2F(M, \Lambda) - 2F(m, \Lambda) - I_1(m) - I_2(m) \} = 0, \quad (12)$$

where the function $F(x, \Lambda)$ is defined in (6),

$$I_i(m) = |e_i H| \left\{ \ln \Gamma(x_i) - \frac{1}{2} \ln(2\pi) + x_i - \frac{1}{2}(2x_i - 1) \ln x_i \right\} \quad (13)$$

and $\Gamma(x)$ is the first order Euler gamma-function [26]. Below we suppose that H is a nonnegative quantity.

The function $F(x, \Lambda)$ vs x monotonically decreases on the interval $x \in (0, \infty)$ (see subsection II.B), and the functions $I_i(x)$ vs x also monotonically decreases on the same interval from $+\infty$ to 0 (see [22]). Hence, for arbitrary fixed values of H, M there exists only one nontrivial solution $m_o(H)$ of equation (12), which is the global minimum point of $\Omega(m; 0, H)$. The quantity $m_o(H)$ is a monotonically increasing function of H and $m_o(H) \rightarrow M$ at $H \rightarrow 0$. It means, that at $G > G_{crit}$ (or for all $M > 0$) the external magnetic field H enhances the discrete chiral symmetry breaking. Note that in the presence of an external magnetic field the parameter m_{vac} from (2) is equal to $m_o(H)$, i.e. depends on the values of H .⁶

⁶ At $G < G_{crit}$, the magnetic catalysis effect of dynamical chiral symmetry breaking (DCSB) takes place. This means that at $H = 0$ the NJL vacuum is chirally symmetric, but an arbitrary small value of the external magnetic field induces DCSB, and fermions acquire a nonzero mass $m_o(H)$ [21, 27] (see also the reviews [28] and recent papers on applications of the magnetic catalysis effect [29]).

Now let us consider the more general case, when $H \neq 0, \mu \neq 0$. Using the methods of [22], it is possible to obtain the thermodynamic potential:

$$\Omega(m; \mu, H) = \Omega(m; 0, H) - \sum_{i=1}^2 \frac{3|e_i|H}{4\pi^2} \sum_{k=0}^{\infty} \alpha_k \theta(\mu - s_{ik}) \left\{ \mu \sqrt{\mu^2 - s_{ik}^2} - s_{ik}^2 \ln \left[\frac{\mu + \sqrt{\mu^2 - s_{ik}^2}}{s_{ik}} \right] \right\}, \quad (14)$$

where $s_{ik} = \sqrt{m^2 + 2|e_i|Hk}$, $\alpha_k = 2 - \delta_{0k}$, and $\Omega(m; 0, H)$ is given in (11). The corresponding stationarity equation at $G > G_{crit}$ looks like:

$$\begin{aligned} \frac{2\pi^2}{3} \frac{\partial}{\partial m} \Omega(m; \mu, H) \equiv m \left\{ 2F(M, \Lambda) - 2F(m, \Lambda) - I_1(m) - I_2(m) + \right. \\ \left. + \sum_{i=1}^2 |e_i|H \sum_{k=0}^{\infty} \alpha_k \theta(\mu - s_{ik}) \ln \left[\frac{\mu + \sqrt{\mu^2 - s_{ik}^2}}{s_{ik}} \right] \right\} = 0. \end{aligned} \quad (15)$$

It is useful to divide the (H, μ) -plane into regions ω_k , which have the following form:

$$(H, \mu) = \bigcup_{k=0}^{\infty} \omega_k, \quad \text{where} \quad \omega_k = \{(H, \mu) : 2|e_1|Hk \leq \mu^2 \leq 2|e_1|H(k+1)\}. \quad (16)$$

Then, the construction of the phase portrait is greatly simplified, since for each region ω_n from (16) only the terms with $k = 0, \dots, n$ from the e_1 -sums in (14)-(15) are nonvanishing, and only the terms corresponding to $k = 0, \dots, 2n+1$ from the e_2 -sums in these expressions are nonzero as well. Hence, it is possible to perform step-by-step the numerical investigation of the phase structure of the NJL model in the regions ω_0, ω_1 , etc. In particular, we have considered in detail the phase structure for the same three values of model parameter $M = M_1, M_2, M_3$ as in the paper [13].

In the case $M = M_1$, we have studied the phase structure only in the region ω_0 (16). The phase portrait in terms of $(H, \mu) \in \omega_0$ is presented schematically on Fig. 2, where $L = \{(H, \mu) : \mu = m_o(H)\}$ denotes the upper estimate of the value μ_c from (2), i.e. $m_{vac} = m_o(H)$. Below L the (H, μ) -plane is divided into four domains A, B, C and C_1 . The solid and dashed lines on this figure correspond to the critical curves of second and first order phase transitions, respectively. The points from the region A correspond to the massless, chirally invariant and dense phase. In region B there is the chirally noninvariant phase with zero baryon density. The one-quark excitations of the vacuum phase B have a mass equal to $m_o(H)$, which depends on H and has no μ -dependency (the phase B can be identified with the real QCD vacuum without baryons). On Fig. 2 one can see also two another massive, i.e. chirally noninvariant, dense phases C and C_1 as well as three tricritical points α, β, γ . Their coordinates are $\sqrt{2|e|H\alpha}/\Lambda = 0.85$, $\mu_\alpha/\Lambda = 0.45$, $\sqrt{2|e|H\beta}/\Lambda = 0.82$, $\mu_\beta/\Lambda = 0.47$ and $\sqrt{2|e|H\gamma}/\Lambda = 0.67$, $\mu_\gamma/\Lambda = 0.48$.

Now let us draw special attention to the boundaries l_{AB} and l_{BC} of the vacuum phase B. The boundary l_{AB} between phases A and B is the graph of the function $\mu = \mu_c(H)$, where $\mu_c(H)$ is the solution of the equation

$$\Omega(0; \mu, H) = \Omega(m_o(H); \mu, H), \quad (17)$$

i.e. on this curve the order parameter – the global minimum point of the thermodynamic potential – changes by a jump its value (from $m_o(H)$ to 0), and a first order phase transition takes place. It turns out that the critical line l_{AB} as a whole is arranged inside the ω_0 -region. However, the boundary l_{BC} , which is also a critical line of first order phase transitions, where the order parameter changes by a jump from $m_o(H)$ to some another definite nonzero value $m_1(H)$, is stretched through the regions $\omega_0, \dots, \omega_{100}$ at least. Evidently, it tends to the point M_1 (see Fig. 2), but to prove this hypothesis, we would need more calculational capacities. The curve l_{BC} is the graph of the function $\mu = \tilde{\mu}_c(H)$, where $\tilde{\mu}_c(H)$ is defined by the equation

$$\Omega(m_1(H); \mu, H) = \Omega(m_o(H); \mu, H). \quad (18)$$

At $M = M_{2,3}$ and $H \neq 0$ the phase portraits at $\mu < m_o(H)$ is qualitatively the same as in the case $H = 0$: there are two phases A and B in this region, and the boundary l_{AB} is the curve of first order phase transitions.

IV. QUARK DROPLET STABILITY AT $H \neq 0$

In the previous section the phase structure of the NJL model was investigated at $H \neq 0$ and for three particular values of the model parameter M : $M = M_1 \equiv 0.48\Lambda$, $M = M_2 \equiv 0.67\Lambda$ and $M = M_3 \equiv 0.88\Lambda$. Now, putting special attention just to these values of M , let us discuss the quark matter stability problem in the presence of an external magnetic field. Stability means that for fixed H there is the critical value μ_c of the chemical potential at which the vacuum and dense quark phases coexist, i.e. at μ_c a first order phase transition occurs. In this case, the value μ_c must satisfy the stability relations (2).

A. Magnetic catalysis of quark matter stability

First of all, let us draw our attention to the case $M = M_1$. The corresponding phase portrait is shown on Fig. 2, where points $(H, \mu = m_o(H))$ are located on the curve L, and $m_o(H)$ is the global minimum point of Ω at $\mu = 0$. Hence, $m_o(H)$ is the mass of the one-quark excitations of the empty vacuum phase B and must be identified with m_{vac} from (2). The boundary of this phase is the first order phase transition line which consists of two curves l_{AB} and l_{BC} and lies below the line L.⁷ So, at $H > H_\alpha$, there is coexistence of the zero density vacuum phase B with the dense phase A on the curve l_{AB} (the one-quark excitations of the phase A have zero mass). Now, if one puts $m_{dense} \equiv 0$, then $\mu_c \equiv \mu_c(H)$ satisfies the stability constraints (2), where $\mu_c(H)$ is defined by the equation (17). Hence, at $H > H_\alpha$, multiquark droplets which are schematically the bags of massless quarks, are allowed to exist in the vacuum.

At $H < H_\alpha$ and in the case $(H, \mu = \tilde{\mu}_c(H)) \in l_{BC}$, where $\tilde{\mu}_c(H)$ is defined by the equation (18), the function $\Omega(m; \mu, H)$ has two different global minimum points: the first one is located at $m = m_o(H)$, the second one at some definite nonzero point $m_1(H)$. So, at $H < H_\alpha$ there is a coexistence of the vacuum phase B with the dense phase C on the curve l_{BC} (the value $m_1(H)$ is just the mass of the one-quark excitations of the ground state of the phase C). Now, if one puts $m_{dense} \equiv m_1(H)$, $m_{vac} \equiv m_o(H)$, then $\mu_c \equiv \tilde{\mu}_c(H)$ satisfies the stability constraints (2). This conclusion can be obtained from a detailed analysis of the stationarity equation (15). At $\Lambda = 650$ MeV it is also confirmed by numerical calculations depicted on Fig. 3 showing graphs of the functions m_{vac} , μ_c and m_{dense} vs H corresponding at $\sqrt{2}|e|\bar{H}/\Lambda < 0.85$ to those points of the line l_{BC} , which lie inside the region ω_0 . So, at $H < H_\alpha$ stable quark droplets composed of *massive* quarks are allowed to exist in the NJL model.

Recall, that in the case $M = M_1$ the stability of quark droplets is absent at $H = 0$ (see Section II as well as [13]). However, as we have proved, in the presence of an external magnetic field there might exist stable quark droplets, i.e. an external magnetic field promotes the stabilization of quark matter. Moreover, it follows from our consideration that at $H \neq 0$, in contrast to the case $H = 0$, stable multiquark clumps might be composed both from massless quarks (at $H > H_\alpha$), and *massive* ones (at $H < H_\alpha$). These are the main results of our paper.

Remark, that usually the value of the cutoff parameter Λ varies in the interval $(0.5 \div 1)$ GeV. In this case, the magnitude of H_α can be estimated to be of the order 10^{19} Gauss.⁸ According to modern estimates, the strength of the magnetic field at the surface of neutron stars can reach 10^{15} Gauss, and at the star core it may be as large as 10^{18} Gauss [30]. Hence, for $M = M_1$ as well as for some neighbouring values of M , the NJL model may predict the existence of stable multiquark clumps, composed of *massive* quarks, inside neutron stars.

Note finally, that for arbitrary values of $M \in (\tilde{M}, M_c)$ (see Fig. 1), quark droplet stabilization is induced by the external magnetic field in the same manner, as in the case $M = M_1$.

At $M = M_{2,3}$ and in the presence of an external magnetic field, the zero density vacuum phase B ($m_{vac} = m_o(H)$) might coexist only with the dense phase A ($m_{dense} = 0$) at $\mu = \mu_c$. Critical values μ_c of the chemical potential in these cases are depicted on Figs 4, 5 for $M = M_2$ and $M = M_3$, respectively. It follows from these figures that the stability constraints (2) are also fulfilled, so the stable quark droplets are allowed to exist. As in the case $H = 0$ (see [13]), at $M = M_{2,3}$ and $H \neq 0$ the finite lumps of stable quark matter are schematically the bags of massless quarks only.

The phase structure of the NJL model at $H \neq 0$ and $\mu \neq 0$ was studied in [21, 22], but for qualitatively other values of model parameters. In particular, the stability problem of multiquark clumps was not considered in these articles. Thus, in view of our present results, let us now discuss it in these cases, too. In [21], in connection with the magnetic catalysis effect, the influence of an external magnetic field on the dense NJL model was investigated at $G < G_{crit}$. It turns out that at sufficiently small values of μ and arbitrary values of H the external magnetic field induces the rearrangement of the model vacuum: at $H = 0$ it is chirally invariant, but the chiral symmetry breaking is induced at $H \neq 0$ (magnetic catalysis effect [27, 28]). Due to the zero baryonic density it is the vacuum phase B. At larger values of μ the dense chirally symmetric phase A with massless quarks is realized. As shown in [21], a first order phase transition occurs between the A and B phases. Now one can easily check that the stability condition (2) of the present paper is fulfilled at $G < G_{crit}$, $H \neq 0$, too. Hence, in this case stable droplets, composed from massless quarks, are allowed to exist. These droplets are surrounded by the chirally noninvariant vacuum. In papers [22] the parameter M was fixed in such a way, that $M < \tilde{M} < M_c$ (see Fig. 1 of the present paper), i.e. at $H = 0$ the quark matter is not stable in this case. Then, a rather complicated phase portrait was shown to exist in the (H, μ) -plane. A more detailed analysis results in the statement that the stability constraints (2) can be solved for these values of model parameters as well (in a similar way, as in the case $M = M_1$).

Hence, at $H \neq 0$, in contrast to the $H = 0$ case, stable quark droplets are allowed to exist in the framework of NJL model (1) for all values of the coupling constant $G > 0$: at $G < G_{bag}$ this property is induced by an external magnetic field, whereas at $G > G_{bag}$ it is an own property of the model.

⁷ On Fig. 2 there is another critical line, where 1st order phase transitions take place as well. It is the line which connects tricritical points β and γ . However, it is the boundary between two dense phases, and no one of them might be identified with the physical vacuum.

⁸ This estimate can be obtained from the known relation $m_e^2/|e| = 4.4 \times 10^{13}$ Gauss (where m_e and e are the mass and the electric charge of an electron, respectively), which permits to connect $(\text{MeV})^2$ and Gauss units.

B. Energetical approach to the stability problem

It might seem that at $M = M_{2,3}$ and $H \neq 0$ the solution of the quark matter stability problem is qualitatively the same as at $H = 0$. However, there are some peculiarities which can be found by an energetical consideration of this problem. Analogously to the case at $H = 0$ [13], let us define the energy density ε at $H \neq 0$ by the following relation

$$\varepsilon = \Omega(m; \mu, H) - \Omega(m_o(H); 0, H) + N_c \mu \rho, \quad (19)$$

where $\rho = -\partial\Omega/(N_c\partial\mu)$ is the baryonic density and $m_o(H)$ – the dynamical quark mass in the vacuum phase B. A very important physical quantity is ε/ρ – the energy per one baryon. At $H = 0$ the graph of this function vs ρ is presented on Fig. 6 for $M = M_2$ and $\Lambda = 600$ MeV (see also [13]). The absolute minimum of this function lies at the point $\rho = \rho_{drop}(M_2) \equiv 2.86\rho_0$ which is equal to the baryonic density inside stable quark droplets. At the point $\rho = 0$ there is a local minimum of ε/ρ , corresponding to the real empty physical QCD-vacuum, i.e. to the phase B. Due to the configuration of the curve ε/ρ vs ρ at $H = 0$, one can conclude that the vacuum, i.e. the state with $\rho = 0$, is a metastable one. It means that not for all values of baryon density fluctuations the system will pass automatically to the mixed phase with stable quark droplet formation. Indeed, the curve ε/ρ has a local maximum at some point ρ_{max} . So, if a quark droplet with $\rho < \rho_{max}$ is created in the vacuum, it is energetically favorable for the system to reach zero density, i.e. to let tend the volume of the quark droplet to infinity thus pushing out all quarks from the system. As a result, one gets the initial empty vacuum. However, if a quark droplet with $\rho > \rho_{max}$ is formed, then it will be a stable one, since in this case the droplet simply changes its volume in order to reach the most energetically favorable density $\rho_{drop}(M_2)$.

The situation is essentially different in the presence of an external magnetic field. In this case the graph of the function ε/ρ has a local maximum in the point $\rho = 0$ (see Fig. 6), so at arbitrary small baryonic density fluctuations the stable quark droplets will appear in the system. A similar property of the function ε/ρ is observed for the case $M = M_1$ (see Fig. 7) as well as for other values of the parameter M . Hence, the external magnetic field promotes the creation of the stable quark matter.

Let us denote the global minimum point of the function ε/ρ as ρ_{drop} (this value is the baryonic density inside the stable quark droplets). Then, the binding energy per quark E_B inside stable droplets is given by the relation

$$E_B \equiv - \left(\frac{\varepsilon}{N_c \rho} \Big|_{\rho=\rho_{drop}} - m_{vac} \right) = m_{vac} - \mu_c, \quad (20)$$

where $m_{vac} = m_o(H)$. Of course, both quantities E_B and ρ_{drop} depend on H . Their graphs are presented on Fig. 8 and Fig. 9, respectively. Recall, that in the case $M = M_2$ the quark binding energy E_B is about 20 MeV at $H = 0$ (see [13]). Now one can easily see from Fig. 8 that at nonzero H this quantity might be several times larger, than at $H = 0$. So, the external magnetic field enhances the stability of quark matter.

Note that on Fig. 9 the density curve corresponding to the case $M = M_2$ has a cusp at the point $\sqrt{2|e|H}/\Lambda \approx 1$. The matter is that in the chirally symmetric phase such quantities, as particle density, magnetization etc, have a discontinuity of their derivatives over μ and H , when a new Landau level is switched on [22]. In the case under consideration in addition to the lowest Landau level the first one for the d -quarks is brought into account just at the point $\sqrt{2|e|H}/\Lambda \approx 1$, so the cusp for the density curve appears. (On Fig. 8, a cusp on the curve E_B vs H also appears at the point $\sqrt{2|e|H}/\Lambda \approx 1$ in the case $M = M_2$, which is, however not so sharp. One can easily see a cusp on the curve μ_c vs H of Fig. 4 as well.) Of course, the smaller the lowest value of the H -interval which is taken into account, the more cusps will appear on the density curve as well as on the graphs of other physical quantities. At $H \rightarrow 0$ there are infinitely many cusps on these curves, which results in the experimentally observed oscillating phenomena of corresponding physical quantities (see [22]).

V. SUMMARY AND DISCUSSIONS

In the present paper the role of an external magnetic field H on the formation of stable quark matter droplets inside the chirally noninvariant vacuum is investigated in the framework of the simple NJL model (1). Earlier, the problem of quark matter stability was considered in terms of more general effective theories with four-fermionic interactions as well, but only at $H = 0$ [13, 14, 15]. Thus, we began our study with the case of a vanishing external magnetic field, too. Moreover, we used the same thermodynamic approach for treating multiquark droplets as in [13]. In this case, any multiquark clump is imagined as a droplet of the dense quark phase of the system, which is surrounded by a chirally noninvariant vacuum phase, and there is thermal and chemical equilibrium between both phases. This qualitative picture can be transformed into a quantitative constraints (2) on dynamical quark masses and the chemical potential at which the coexistence of two phases is possible. Moreover, it was shown in [13] that at some values of the physically acceptable model parameters (coupling constants, cutoff parameter, etc) the stability constraints (2) are not fulfilled for the NJL model at $H = 0$. In those cases stable multiquark droplets are forbidden, and the NJL model is not applicable for the description of dense quark matter.

In our paper we have in the first step obtained the general solution of the stability constraints (2) at $H = 0$ for the NJL model considered above. Here, we have considered only the physically interesting case, when $G > G_{crit} = \pi^2/(3N_f\Lambda^2)$, in which case chiral symmetry is spontaneously broken down and quarks acquire a dynamical mass M (at $\mu = 0$). Due to the stationarity equation (5), the quantities M and G (at $G > G_{crit}$) are in a one-to-one correspondence. Thus, we find it convenient to use the value M (instead of G) as a model parameter (if M changes from 0 to ∞ , then G monotonically increases from G_{crit} to ∞). Stable quark matter droplets inside the chirally noninvariant vacuum are then allowed in the model under consideration only at $M > M_c \equiv 0.56\Lambda$, i.e. at $G > G_{bag} \equiv 1.37G_{crit}$, and they are composed of massless quarks. Notice also that in [13] the stability problem was investigated only at three physically admissible sets of parameters: $M = M_1 \equiv 0.48\Lambda$, $\Lambda = 650$ MeV (set 1), $M = M_2 \equiv 0.67\Lambda$, $\Lambda = 600$ MeV (set 2) and $M = M_3 \equiv 0.88\Lambda$, $\Lambda = 570$ MeV (set 3). There it was pointed out that for set 1 stable quark droplets are forbidden, but for sets 2 and 3 stable quark matter is allowed to exist. Of course, these results coincide with our ones at $H = 0$.

Next, we took into account an external magnetic field and studied the problem of quark matter stability in detail at the same values of the parameter M as in ref.[13]. It turns out that at any value of H in the cases $M = M_{2,3}$ stable quark droplets are allowed to exist. Moreover, as at $H = 0$, these multi-quark droplets consist of massless quarks.⁹ The most interesting results were obtained in the case $M = M_1$, where we have found that an external magnetic field induces stable quark matter formation. Indeed, at $H = 0$ multi-quark droplets are not stable in this case. However, at $H \neq 0$ they are allowed to exist. Furthermore, there exists a value of magnetic fields, $H_\alpha \sim 10^{19}$ Gauss, such that at $H < H_\alpha$ quark droplets are formed by *massive* quarks. At $H > H_\alpha$, multi-quark droplets are composed of massless quarks. We have also checked that in a similar way, for other values of M from the interval $M \in (M, M_c)$ (see Fig. 1), stable quark droplet formation is also induced by the external magnetic field.

Note that one can even assert that at $H \neq 0$ stable quark droplets are allowed to exist in the initial NJL model at *arbitrary* values of the coupling constant $G > 0$. At $G > G_{bag}$ it is the native property of this theory even at $H = 0$, but at $G < G_{bag}$ the stability of the quark matter is induced (catalized) by the external magnetic field.

Moreover, using an energetical approach, we have shown that an external magnetic field promotes the creation as well as enhances the stability of quark droplets at $G > G_{bag}$.

Finally, let us add some remarks about the role of color superconductivity (CSC) in the quark matter stability problem considered above. This phenomenon was predicted quite recently in the framework of instanton-induced- as well as NJL-type models with four – fermionic interactions (see the reviews [31] or some of the recent papers [32, 33] and references therein). At zero temperature it is proved that CSC must exist at asymptotically large values of the chemical potential (or baryonic densities), however, the lower boundary for this effect is not established up to now. In particular, in [34] CSC was investigated just in a slightly extended version of the NJL model (1) at neighboring to the set 1 values of parameters, i.e. for $M \approx M_1$ and $\Lambda = 650$ MeV, but with an additional four-quark term included, which is responsible for the interactions in the diquark channel (with a corresponding coupling constant G'). It was realized at $H = 0$, that usually with growing value of μ , the chiral invariance of the system is restored before or just at the CSC phase transition. Moreover, at fixed G , the parameters of the CSC phase transition strongly depend on the value of $\kappa \equiv G'/G$, while the features of the chiral phase transition do not seriously depend on κ . For example, in the NJL model (1) at small values of $\kappa \leq 1/4$ the transition to the CSC phase occurred at $\mu \sim 700$ MeV, but at $\kappa \sim 1$ it is at $\mu \sim 300$ MeV, whereas for wide interval of κ -values (G is fixed) the chiral symmetry is restored at $\mu \sim 300$ MeV [34].

For simplicity, we have put $G' = 0$ in our consideration of the quark matter stability problem at $H \neq 0$. In the light of the above discussion we believe that our results are not qualitatively changed at least for small values of κ , i.e. in the case, where CSC occurs for rather large values of $\mu \sim 700$ MeV [34]. In fact, in this case CSC can not seriously influence stable quark droplet formation which takes place at values of the chemical potential much less than 700 MeV (see the graphs of μ_c on Figs 3-5). At present time there are several indications on the possibility for CSC to take place only at chemical potential values which indeed are outside the scope of our consideration, i.e. at $\mu \gg 300$ MeV: i) the CSC phenomenon was not discovered up to now in heavy ion collision experiments, ii) It was argued in [35] that two-flavor CSC is absent inside quark cores of neutron stars, where baryonic density is several times the density ρ_o of ordinary nuclear matter, iii) In [36] it was shown that due to the color Meissner effect as well as due to the presence of background chromomagnetic fields (gluon condensate) the CSC is shifted to higher densities (chemical potentials), iv) As in the case with ordinary superconductivity, the sufficiently strong external magnetic field prevents a system from CSC as well etc. In these cases the ordinary, i.e. without CSC, quark matter might be formed at rather low densities. To describe such a processes one could use models of the type (1) with zero or very small value of the diquark channel coupling constant G' .

Recently, we have studied already the magnetic catalysis of the quark droplet stabilization in the scope of a simpler one-flavor NJL model [37]. Next, we are planning to consider the quark matter stability problem in more realistic

⁹ It is necessary to note, that for such values of M which are greater, but quite near to M_c (8) (for example, at $M_c < M < 1.01M_c$), there exists a finite interval of the external magnetic field values, where stable quark droplets are formed by massive quarks. But for values of the external magnetic field outside of this interval, stable multi-quark droplets are composed of massless quarks.

NJL models including vector interactions as well as the third flavor of quarks etc.

Acknowledgments

The authors are grateful to Dr. M.L. Nekrasov for help in numerical calculations. This work was supported in part by DFG-project 436 RUS 113/477/0-2.

-
- [1] Z. Fodor and S.D. Katz, Phys. Lett. **B534** (2002) 87; S. Ejiri, C.R. Allton, S.J. Hands et al, hep-lat/0209012; M. D'Elia and M.P. Lombardo, Phys. Rev. **D67** (2003) 014505; F. Karsch and E. Laermann, hep-lat/0305025.
 - [2] J.D. Walecka, Ann. Phys. **83** (1974) 491.
 - [3] J.I. Kapusta, *Finite-temperature field theory*, (Cambridge University Press,1989).
 - [4] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345.
 - [5] D. Ebert and M.K. Volkov, Yad. Fiz. **36** (1982) 1265; Z.Phys. **C16** (1983) 205; D. Ebert and H. Reinhardt, Nucl. Phys. **B271** (1986) 188; D. Ebert, H. Reinhardt and M.K.Volkov, Progr. Part. Nucl. Phys. **33** (1994) 1.
 - [6] D. Ebert, L. Kaschluhn and G. Kastelewicz, Phys. Lett. **B264** (1991) 420.
 - [7] U. Vogl, Z. Phys. **A337** (1990) 191; U. Vogl and W. Weise, Progr. Part. Nucl. Phys. **27** (1991) 195.
 - [8] D. Ebert and L. Kaschluhn, Phys. Lett. **B297** (1992) 367; D. Ebert and T. Jurke, Phys. Rev. **D58** (1998) 034001; L.J. Abu-Raddad et al., Phys. Rev. **C66** (2002) 025206.
 - [9] H. Reinhardt, Phys. Lett. **B244** (1990) 316; N. Ishii, W. Benz and K. Yazaki, Phys. Lett. **B301** (1993) 165; C. Hanhart and S. Krewald, ibid. **344** (1995) 55.
 - [10] T. Hatsuda and T. Kunihiro, Phys. Rept. **247** (1994) 221.
 - [11] W.M. Alberico, F. Giacosa, M. Nardi and C. Ratti, Eur. Phys. J. **A16** (2003) 221.
 - [12] M. Oettel, G. Hellstern, R. Alkhofer and H. Reinhardt, Phys. Rev. **C58** (1998) 2459.
 - [13] M. Buballa, Nucl. Phys. **A611** (1996) 393; M. Buballa and M. Oertel, Nucl. Phys. **A642** (1998) 39.
 - [14] M. Alford, K. Rajagopal and F. Wilczek, Phys. Lett. **B422** (1998) 247.
 - [15] M. Buballa and M. Oertel, Phys. Lett. **B457** (1999) 261; I.N. Mishustin, L.M. Satarov, H. Stucker et al, Phys. Atom. Nucl. **64** (2001) 802; C. Ratti, Europhys. Lett. **61** (2003) 314.
 - [16] A. Chodos, R.L. Jaffe, K. Johnson and C.B. Thorn, Phys. Rev. **D10** (1974) 2599.
 - [17] A.S. Vshivtsev and D.V. Serebryakova, Zh. Eksp. Teor. Fiz. **106** (1994) 35; G.G. Likhachev and A.I. Studenikin, JETP **81** (1995) 419; S. Chakrabarty, Phys. Rev. **D54** (1996) 1306; T.C. Phukon, Phys. Rev. **D62** (2000) 023002; V.R. Khalilov, Phys. Rev. **D65** (2002) 056001; Teor. Mat. Fiz. **133** (2002) 103.
 - [18] T. Tatsumi, Phys. Lett. **B489** (2000) 280; astro-ph/0004062.
 - [19] M. Alford, J. Berges and K. Rajagopal, Nucl. Phys. **B571** (2000) 269.
 - [20] D. Blaschke, D.M. Sedrakian and K.M. Shahabasyan et al, Astrofiz. **44** (2001) 443.
 - [21] K.G. Klimenko, preprint IHEP 98-56, Protvino 1998 [hep-ph/9809218]; A.S. Vshivtsev, M.A. Vdovichenko and K.G. Klimenko, Phys. Atom. Nucl. **63** (2000) 470.
 - [22] D. Ebert et al., Phys. Rev. **D61** (2000) 025005; D. Ebert, M.A. Vdovichenko and K.G. Klimenko, Phys. Atom. Nucl. **64** (2001) 336; D. Ebert and K.G. Klimenko, nucl-th/9911073.
 - [23] J. Madsen, Phys. Rev. **D47** (1993) 5156; E.P. Gilson and R.L. Jaffe, Phys. Rev. Lett. **71** (1993) 332; O. Kiriya and A. Hosaka, Phys. Rev. **D67** (2003) 085010.
 - [24] P. Zhuang, J. Hufner and S.P. Klevansky, Nucl. Phys. **A576** (1994) 525.
 - [25] A.S. Vshivtsev and K.G. Klimenko, JETP Lett. **64** (1996) 338 [hep-ph/9701288]; A.S. Vshivtsev, V.Ch. Zhukovsky and K.G. Klimenko, JETP **84** (1997) 1047.
 - [26] H. Bateman and A. Erdelyi, *Higher Transcendental Functions*, (McGrawHill, New York, 1953).
 - [27] K.G. Klimenko, Theor. Math. Phys. **89** (1991) 1161; **90** (1992) 1; Z. Phys. **C54** (1992) 323; I.V. Krive and S.A. Naftulin, Phys. Rev. **D46** (1992) 2737; V.P. Gusynin, V.A. Miransky and I.A. Shovkovy, Phys. Rev. Lett. **73** (1994) 3499; Phys. Lett. **B349** (1995) 477.
 - [28] V.A. Miransky, Progr. Theor. Phys. Suppl. **123** (1996) 49; A.S. Vshivtsev et al, Phys. Part. Nucl. **29** (1998) 523; V.P. Gusynin, Ukr. J. Phys. **45** (2000) 603; V. de la Incera, hep-ph/0009303.
 - [29] V.C. Zhukovsky et al, JETP Lett. **73** (2001) 121; Theor. Math. Phys. **134** (2003) 254; E.J. Ferrer, V.P. Gusynin and V. de la Incera, cond-mat/0203217; V.A. Miransky, hep-ph/0208180; E. Elizalde, E.J. Ferrer and V. de la Incera, hep-ph/0209324; V.A. Miransky and I.A. Shovkovy, Phys. Rev. **D66** (2002) 045006; V.P. Gusynin, V.A. Miransky and I.A. Shovkovy, hep-ph/0304059.
 - [30] R.C. Duncan, astro-ph/0002442; S. Ghosh et al, astro-ph/0106153.
 - [31] M. Alford, Ann. Rev. Nucl. Part. Sci. **51** (2001) 131; B.O. Kerbikov, hep-ph/0110197; G. Nardulli, Riv. Nuovo Cim. **25N3** (2002) 1.
 - [32] S.M. Molodtsov and G.M. Zinovjev, Mod. Phys. Lett. **A18** (2003) 817; hep-ph/0112075; B.O. Kerbikov, Phys. Atom. Nucl. **65** (2002) 1918; M. Huang, P. Zhuang and W. Chao, Phys. Rev. **D65** (2002) 076012; hep-ph/0110046; E. Nakano, T. Maruyama and T. Tatsumi, hep-ph/0304223.
 - [33] D. Ebert, K.Klimenko and H.Toki, Phys.Rev. **D64** (2001) 014038; D. Ebert et al. Prog. Theor. Phys. **106** (2001) 835; JETP Letters **74** (2001) 523.

- [34] T.M. Schwarz, S.P. Klevansky and G. Papp, Phys. Rev. **C60** (1999) 055205; M. Sadzikowski, Mod. Phys. Lett. **A16** (2001) 1129.
- [35] M. Alford and K. Rajagopal, hep-ph/0204001.
- [36] N.O. Agasian, B.O. Kerbikov and V.I. Shevchenko, Phys. Rept. **320** (1999) 131; G.M. Carter and D. Diakonov, Nucl. Phys. **B582** (2000) 571; D. Ebert et al, Phys. Rev. **D65** (2002) 054024.
- [37] K.G. Klimenko and D. Ebert, *Magnetic catalysis of the quark matter stability in the Nambu – Jona-Lasinio model*, preprint IFVE 2003-5, Protvino (2003) [in Russian].

Figure Captions

Fig. 1. Schematic phase portrait of the NJL model at nonzero μ and for arbitrary values of the quark mass M at $\mu = 0$ (the case $G > G_{crit}$). Phases B and C are massive nonsymmetric phases, A is a chirally symmetric and massless phase. In the phase B the quark mass is equal to M . In the phase B the particle density in the ground state is equal to zero, whereas in phases A, C it is nonzero. Solid and dashed lines represent critical curves of second and first order phase transitions, respectively; α and β denote tricritical points. In particular, the curve l_{BC} has the following form: $l_{BC} = \{(M, \mu) : \mu = M\}$, the curve l_{AB} is defined by the equation (7), $\tilde{M} = 0.31\Lambda$, $M_c = 0.56\Lambda$. Only the values $\mu_c(M)$ (see (8)) of the chemical potential, corresponding to points of the critical line l_{AB} , obey the stability constraints (2).

Fig. 2. Schematic (H, μ) -phase portrait of the NJL model at $M = M_1 \equiv 0.48\Lambda$ ($\tilde{M} < M_1 < M_c$), where A is the chirally symmetric massless phase, B, C and C_1 are chirally noninvariant massive phases. Ground state baryonic density is zero in the phase B, and nonzero in phases A, C, C_1 . Dashed and solid lines are critical curves of the first- and second order phase transitions, respectively. (l_{AB} is defined by the equation (17)) The curve $L = \{(H, \mu) : \mu = m_o(H)\}$, where $m_o(H) \equiv m_{vac}$ is the quark mass at $\mu = 0$. $\mu_1 \equiv 326$ MeV is such a value of the chemical potential that the point (M_1, μ_1) lies on the curve $\widehat{\alpha\beta}$ of Fig. 1. The external magnetic field values, corresponding to the tricritical points α, β, γ are the following: $\sqrt{2|e|H_\alpha}/\Lambda = 0.85$, $\sqrt{2|e|H_\beta}/\Lambda = 0.82$ and $\sqrt{2|e|H_\gamma}/\Lambda = 0.67$. Only the points of the critical lines l_{AB} and l_{BC} obey the stability constraints (2).

Fig. 3. The behaviour of quantities $m_{vac} \equiv m_o(H)$, μ_c and m_{dense} vs H which characterize the coexistence of the phase B with A and C ones in the case $M = M_1$ and $\Lambda = 650$ MeV. If $\sqrt{2|e|H}/\Lambda > 0.85$ these parameters describe the first order phase transition (curve l_{AB}) between A and B (in this case $m_{dense} \equiv 0$ and $\mu_c \equiv \mu_c(H)$, where $\mu_c(H)$ is the solution of the equation (17)), if $\sqrt{2|e|H}/\Lambda < 0.85$ – the phase transition between B and C (in this case $m_{dense} \equiv m_1(H)$ and $\mu_c \equiv \tilde{\mu}_c(H)$, where $\tilde{\mu}_c(H)$ is defined by the equation (18)). Only the part of l_{BC} , lying in the region ω_0 is considered. These values of m_{vac} , μ_c and m_{dense} are the solutions of the stability constraints (2).

Fig. 4. The same as on Fig. 3, but for $M = M_2$ and $\Lambda = 600$ MeV (in this case $m_{dense} \equiv 0$). The dotted horizontal line corresponds to the critical chemical potential value μ_c at zero external magnetic field. For these values of m_{vac} , μ_c and m_{dense} the stability constraints (2) are fulfilled.

Fig. 5. The same as on Fig. 3, but for $M = M_3$ and $\Lambda = 570$ MeV (in this case $m_{dense} \equiv 0$ as well). The dotted horizontal line corresponds to the critical chemical potential value μ_c at zero external magnetic field. For these values of m_{vac} , μ_c and m_{dense} the stability constraints (2) are fulfilled.

Fig. 6. The energy per baryon in the NJL model (1) at $M = M_2$, $\Lambda = 600$ MeV for $H = 0$ and $\sqrt{2|e|H}/\Lambda = 1.04$. The solid line corresponds to the massive solutions of the stationarity equation, the pointed line to the massless one.

Fig. 7. The energy per baryon in the NJL model (1) at $M = M_1$, $\Lambda = 650$ MeV for $\sqrt{2|e|H}/\Lambda = 0.82$ and $\sqrt{2|e|H}/\Lambda = 1.04$. The solid line corresponds to the massive solutions of the stationarity equation, the pointed line to the massless one.

Fig. 8. The binding energy per quark for a stable droplet in the NJL model (1) vs external magnetic field in two cases: $M = M_1$, $\Lambda = 650$ MeV (solid line) and $M = M_2$, $\Lambda = 600$ MeV (pointed line).

Fig. 9. Baryon density ρ_{drop} inside a stable quark droplet in the NJL model (1) as a function of external magnetic field. The solid line corresponds to the case $M = M_1$, $\Lambda = 650$ MeV, the pointed one to the case $M = M_2$, $\Lambda = 600$ MeV.




FIG. 1:




FIG. 2:




FIG. 3:




FIG. 4:




FIG. 5:




FIG. 6:




FIG. 7:




FIG. 8:




FIG. 9: